

Winchester Surgeons

“Putting Patient Care First”



Damian Szewczyk, M.D, FACS
Troy Hixson, M.D, FACS

83 Memorial Drive · Winchester, TN 37398
Phone: 931-967-3966 Fax: 931-962-0373
www.winsurgeons.net

Post-operative instructions following inguinal hernia surgery

Diet: You can eat a regular diet without any restrictions after surgery. On the first day eat small meals and drink plenty of liquids (8-10 glasses). Nausea and vomiting from anesthesia is not uncommon for 24 hours after surgery. Do not take any alcoholic drinks while taking pain killers prescribed by your doctor.

Activity: You can walk during the day to help your body return to normal. You can walk up stairs. Do not lift more than 10-15 pounds for 4-6 weeks from surgery. Do not drive if you are taking narcotic medicine (Norco/Lortab, Percocet, Demerol, Tylenol #3 etc.) or within 24 hours afterward. Do not return to exercise or strenuous activity until you are seen in follow-up. You can return to work when you feel comfortable, and when you can perform your job safely and at the level you and your employer expect.

Pain and discomfort: You have been given a prescription for a narcotic (Norco/Lortab, Percocet, Tylenol#3, Demerol or Dilaudid). Call the office if you experience itching or a rash and stop the medication. If you don't need as much pain medicine, you can take two extra strength Tylenol every 6 hours. All narcotics will make you constipated. Drink plenty of liquids. If you do not have a bowel movement in 2-3 days then take one of the following by mouth: Miralax (1 serving-2 times a day) or Metamucil (1 Tablespoon in 16oz of water or juice-2 times a day) or Mineral Oil (2 tablespoons-2 times a day) or Milk of Magnesia (2 tablespoons-2 times a day).

Incisional care: You may experience some swelling and bluish discoloration at the incision site and in men at the scrotum and penis. This is normal. You may use an ice pack for 24-48 hours to alleviate the swelling and discomfort. You may brace your incision with a small pillow when coughing and/or sneezing. Your incisions are covered with skin glue-you may shower after you go home in 24 hrs. Do not soak in a tub or swimming pool. The sutures under the skin will dissolve 3-4 weeks after surgery.

Call our office if you have: Temperature over 101.5F, redness or swelling at the incision site, foul smelling drainage from the incision, increased tenderness or soreness at the wound or the wound edges are no longer together.

Follow up appointment: Please call our office at your convenience to schedule a follow-up visit 7-10 days after your surgery.